

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

Publication number:

0 507 317 A2

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 92105776.6

(51)

Int. Cl.5: **C07C 279/26, A61K 31/155**

(22) Date of filing: 03.04.92

(30) Priority: 05.04.91 JP 73202/91
 04.09.91 JP 224306/91
 19.06.91 JP 147644/91

(43) Date of publication of application:
 07.10.92 Bulletin 92/41

(84) Designated Contracting States:
**AT BE CH DE DK ES FR GB GR IT LI LU MC
 NL PT SE**

(71) Applicant: OTSUKA PHARMACEUTICAL CO.,
 LTD.
 9, Kandatsukasacho 2-chome
 Chiyoda-ku Tokyo 101(JP)

(72) Inventor: Ishikawa, Hiroshi
 18-14, Seifu-cho
 Otsu-shi, Shiga 520-02(JP)
 Inventor: Yasumura, Koichi


8-11-401, Uchidehama
 Otsu-shi, Shiga 520(JP)
 Inventor: Tsubouchi, Hidetsugu
 2-6-608, Nionohama 2-chome
 Otsu-shi, Shiga 520(JP)
 Inventor: Higuchi, Yukio
 8-17, Chodo 2-chome
 Higashiosaka-shi, Osaka 577(JP)
 Inventor: Tamaoka, Hisashi
 5-32, Aza-ekogawa, Shinkirai, Kitajima-cho
 Itano-gun, Tokushima 771-02(JP)

(74)

Representative: Hansen, Bernd, Dr.
 Dipl.-Chem. et al
 Hoffmann, Eitle & Partner Patent- und
 Rechtsanwälte Arabellastrasse 4 Postfach
 81 04 20
 W-8000 München 81(DE)


(54) **Biguanide derivatives, manufacturing method thereof, and disinfectants containing the derivatives.**

(57) The invention presents a biguanide derivative or its salt expressed by a formula:



(where R¹ and R² are as defined in Specification), or formula:

EP 0 507 317 A2



(2)

(where A and R³ is as defined in specification). This biguanide derivative or its salt is preferably used as the effective amount of a disinfectant for humans, animals, medical appliances, etc.

BACKGROUND OF THE INVENTION


The invention relates to novel biguanide derivatives used in disinfection of humans, animals, medical appliances, and others, a method of manufacturing the same, and disinfectants containing such derivatives.

It is known that various guanidine derivatives possess bactericidal actions, and biguanide derivatives of 1,4-dimethylene cyclohexane type are disclosed in the British Patent Specification No. 1,095,902.

SUMMARY OF THE INVENTION

It is a primary object of this invention to present novel biguanide derivatives or their salts possessing a higher activity than the conventional biguanide derivatives or their salts, and disinfectants containing such derivatives or their salts.

To achieve the above object, a monobiguanide derivative of the invention is a biguanide derivative or its salt, expressed by a general formula:



(where R¹ represents 3,4-dichlorobenzyl group, 4-chlorophenyl group, 3,4-dichlorophenyl group, benzyl group or 4-chlorobenzyl group, and R² represents octyl group, 3,4-dichlorobenzyl group, dodecyl group, decyl group, 3-trifluoromethylphenyl group, 4-bromophenyl group, 4-iodophenyl group, 2,4-dichlorophenyl group, 3,4-dichlorophenyl group, 2,3,4-trichlorophenyl group, 3,4-dimethylphenyl group, 3,4-methylenedioxyphenyl group, 4-*t*-butylphenyl group, 4-ethylthiophenyl group, 1,1,3,3-tetramethylbutyl group, hexyl group, 2-ethoxyethyl group, 2-(2-hydroxyethoxy)ethyl group, 3-diethylaminopropyl group, 3-(2-ethylhexyloxy)-propyl group, (3-isopropoxy)propyl group, (2-diethylamino)-ethyl group, (3-butyl)-propyl group, 3-(*n*-butylamino)propyl group, cyclohexylmethyl group, 3-trifluoromethylphenyl group, 4-ethylthiophenyl group, 4-chlorobenzyl group, 2,4-dichlorobenzyl group, 4-acetylaminothiophenyl group, 3,4-methylenedioxyphenyl group, 3,4-methylenedioxybenzyl group, octyl group, 4-chlorobenzyl group, decyl group, dodecyl group, isobutyl group, 3,4-dichlorophenyl group, or hexyl group), and the derivative being N¹-(3,4-dichlorobenzyl)-N⁵-octyl-biguanide,

N¹-(4-chlorophenyl)-N⁵-(3,4-dichlorobenzyl)-biguanide,
 N¹-(3,4-dichlorophenyl)-N⁵-octyl-biguanide,
 N¹-benzyl-N⁵-dodecyl-biguanide,
 N¹-benzyl-N⁵-decyl-biguanide,
 N¹-(3,4-dichlorobenzyl)-N⁵-(3-trifluoromethylphenyl)-biguanide,
 N¹-(3,4-dichlorobenzyl)-N⁵-(4-bromophenyl)-biguanide,
 N¹-(3,4-dichlorobenzyl)-N⁵-(4-iodophenyl)-biguanide,
 N¹-(3,4-dichlorobenzyl)-N⁵-(2,4-dichlorophenyl)-biguanide,
 N¹-(3,4-dichlorobenzyl)-N⁵-(3,4-dichlorophenyl)-biguanide,
 N¹-(3,4-dichlorobenzyl)-N⁵-(2,3,4-trichlorophenyl)-biguanide,
 N¹-(3,4-dichlorobenzyl)-N⁵-(3,4-dimethylphenyl)-biguanide,
 N¹-(3,4-dichlorobenzyl)-N⁵-(3,4-methylenedioxyphenyl)-biguanide,
 N¹-(3,4-dichlorobenzyl)-N⁵-(4-*t*-butylphenyl)-biguanide,
 N¹-(3,4-dichlorobenzyl)-N⁵-(4-ethylthiophenyl)-biguanide,
 N¹-(3,4-dichlorobenzyl)-N⁵-(*n*-decyl)-biguanide,
 N¹-(3,4-dichlorobenzyl)-N⁵-(1,1,3,3-tetramethylbutyl)-biguanide,
 N¹-(3,4-dichlorobenzyl)-N⁵-hexyl-biguanide,
 N¹-(3,4-dichlorobenzyl)-N⁵-(2-ethoxyethyl)-biguanide,
 N¹-(3,4-dichlorobenzyl)-N⁵-[2-(2-hydroxyethoxy)ethyl]-biguanide,
 N¹-(3,4-dichlorobenzyl)-N⁵-(3-diethylaminopropyl)-biguanide,
 N¹-(3,4-dichlorobenzyl)-N⁵-[3-(2-ethylhexyloxy)-propyl]-biguanide,
 N¹-(3,4-dichlorobenzyl)-N⁵-[(3-isopropoxy)propyl]-biguanide,
 N¹-(3,4-dichlorobenzyl)-N⁵-[(2-diethylamino)-ethyl]-biguanide,
 N¹-(3,4-dichlorobenzyl)-N⁵-[(3-butyl)-propyl]-biguanide,

5
10

20

(2):



35

bigu

40

DE

45

50


55

Such acid-addition salt itself exhibits a high bactericidal effect or antibacterial effect, and is also useful for refining or isolation of free base. Meanwhile, the acid-addition salt is preferred to be acceptable pharmaceutically.


The monobiguanide derivative (1) of this invention may also form a stable coordination compound with a metal salt. Such coordination compound also presents a bactericidal effect. The coordination compound is obtained by reaction of its derivative or its acid-addition salt with a specified quantity of metal salt, preferably a heavy metal salt such as V, Cr, Mn, Co, Ni, Cu, Zn, Pd, Re and Os.

Next is explained a manufacturing method of the monobiguanide derivative (1) of the invention.

10



15



20


25 (where R¹ and R² are same as defined above.)

As shown in the reaction formula above, the monobiguanide derivative (1) of the invention is obtained by reaction of the amine expressed by Formula (a) with N¹-cyanoguanidine compound expressed by Formula (b). At this time, the amine (a) is used in about equimolecular amounts of the compound (b). The amine (a) may be used in a form of proper acid-addition salt such as hydrochloric acid salt.


30 The both components (a) and (b) are caused to react by heating in the presence or absence of inert solvent, such as 2-ethoxyethanol, 2-methoxyethanol and o-dichlorobenzene. If the product is an acid-addition salt, it may be changed into a form of free base by causing an alkali such as sodium hydroxide to act thereon, or may be changed into other acid-addition salt by ion exchange or other process.

35 The starting material expressed by Formula (b) may be manufactured, for example, in the following reaction formula.

40



45




50

(where M represents an alkali metal, and R² is same as defined above.)

That is, the amine expressed by Formula (c) or its acid-addition salt (hydrochloride, etc.) is caused to react with the alkali metal salt (d) of dicyanamide (for example, sodium dicyanamide, potassium dicyanamide) in an inert solvent to obtain the material.

55 The monobiguanide derivative (1) of this invention may be also manufactured in the following reaction formula.



(where R¹ and R² represent the same as above.)

This reaction is performed nearly in the same condition as in the above reaction.

The bis-biguanide derivative (2) of this invention is described below. Examples of halogen atom represented by R³ in Formula (3) may include chlorine, bromine, iodine and fluorine.

Examples of lower alkyl group include straight-chain or branched alkyl group having 1 to 6 carbon atoms, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-amyl, n-hexyl group, and the like.

Examples of halogen substituted lower alkyl group include halogen substituted lower alkyl group having 1 to 6 carbon atoms and replaced by 1 to 3 halogen atoms, such as monochloromethyl, monobromomethyl, monofluoromethyl, monoiodomethyl, dichloromethyl, dibromomethyl, difluoromethyl, diiodomethyl, trichloromethyl, tribromomethyl, trifluoromethyl, triiodomethyl, 3-chloropropyl, 3-fluoropropyl, 2,3-dichloropropyl, 3,3,3-trichloropropyl, 3-chloro-2-methylpropyl, 4-chlorobutyl, 4-fluorobutyl, 5-chloropentyl, 6-chlorohexyl, 6-fluorohexyl group, and the like.


Examples of lower alkylthio group include lower alkylthio group having straight-chain or branched alkyl moiety having 1 to 6 carbon atoms, such as methylthio, ethylthio, n-propylthio, isopropylthio, n-butylthio, isobutylthio, tert-butylthio, n-amylthio, n-hexylthio group, and the like.











Examples of phenyl lower alkyl group which may possess a halogen atom on a phenyl group as a substituent may include phenyl lower alkyl group possessing one to three phenyl groups which may have one to three halogen atoms, and having 1 to 6 carbon atoms in the alkyl moiety, such as benzyl, α -phenethyl, β -phenethyl, 3-phenylpropyl group, benzhydryl, trithyl, 4-chlorophenylmethyl, 3-chlorophenylmethyl, 2-chlorophenylmethyl, 3,4-dichlorophenylmethyl, 4-fluorophenylmethyl, 3,4-difluorophenylmethyl, 4-chlorophenylethyl, 3,4-dichlorophenylethyl group, and the like.









Examples of alkyl group used as substituent R³ are straight-chain or branched alkyl groups having 6 to 16 carbon atoms, specifically including n-hexyl, n-heptyl, n-octyl, n-nonyl, n-decyl, n-undecyl, n-tetradecyl, n-hexadecyl, 2-ethyl-1-hexyl, 2-ethyl-1-heptyl, 2-heptyl, 2-octyl, and 1,1,3,3-tetramethylbutyl group.

Examples of lower alkylene group include alkylene group having 1 to 6 carbon atoms, such as methylene, methylenemethylene, ethylene, dimethylenemethylene, trimethylene, 1-methyl-1-trimethyl, 2-methyl-trimethylene, 2,2-dimethyltrimethylene, tetramethylene, pentamethylene, hexamethylene, and the like.

Practical examples of the bis-biguanide derivative (2) of the invention are shown in the table below.



R ³	A
	—CH ₂ —
	"
	"
	"
	"
	"
	"
	"
	"
	"

	R^3	A
5		$-\text{CH}_2-$
10		"
15		"
20		$-\text{CH}_2\text{CH}_2-$
25		"
30		$-\text{CH}_2-$
35		"
40		$-(\text{CH}_2)_3-$
45	$\text{H}_3\text{C}-$	$-\text{CH}_2-$
50	$(\text{H}_3\text{C})_3\text{C}-$	"

The bis-biguanide derivative (2) of this invention is a basic compound, and forms, same as the monobiguanide derivative (1), a corresponding acid-addition salt by reaction with various organic acids or inorganic acids.

The acid-addition salt of the bis-biguanide derivative (2) itself, as well as its derivative (2), exhibits a high bactericidal effect or antibacterial effect, and it is also useful for refining or isolation of free base. The acid-addition salt is preferred to be acceptable pharmaceutically. The bis-biguanide derivative (2) of the

invention, same as the monobiguanide derivative (1), may also form a stable coordination compound with a metal salt, and such coordination compound also presents an excellent bactericidal effect.


The bis-biguanide derivative (2) of the invention is manufactured, for example, in the following reaction formula.

5

10

15

20



(where A and R³ represent same as defined above.)

As shown in the reaction formula above, the bis-biguanide derivative (2) of this invention is obtained by the reaction of 1,1'-[1,3-cyclohexanebis(alkylene)]-bis(3-cyanoguanidine) shown in Formula (e) with the amine shown in Formula (f). At this time, the amine (f) is used by about twice the mol equivalent of the compound (e). The amine (f) may be in a form of proper acid-addition salt such as hydrochloride.


The both components (e) and (f) are caused to react with each other by heating in the presence or absence of an inert solvent (e.g. 2-ethoxyethanol, 2-methoxyethanol, o-dichlorobenzene). If the reaction product is an acid-addition salt, it may be changed into a form of a free base by causing an alkali such as sodium hydroxide to act thereon, or changed into other acid-addition salt by ion exchange or other process.

The starting material expressed by Formula (e) may be manufactured, for example, in the following reaction formula.

35

40

45




(where M denotes an alkali metal, and A is same as defined above.)

It is obtained, accordingly, by reaction of the diamine shown in Formula (g) or its acid-addition salt (hydrochloride, etc.) with the alkali metal salt (h) of dicyanamide (e.g. sodium dicyanamide, potassium dicyanamide) in an inert solvent.

The bis-biguanide derivative (2) of this invention may be also manufactured in the following reaction formula.

55



(where A and R³ represent the same as above.)

As shown in the above reaction formula, the bis-biguanide derivative (2) is obtained by the reaction between the diamine expressed by Formula (i) or its acid-addition salt (hydrochloride, etc.) and the cyanoguanidine compound expressed by Formula (j), in the presence or absence of an inert solvent. The reaction is conducted under heating. The cyanoguanidine compound (j) is used at a rate of about twice the mol equivalent of the diamine (i).

The biguanide derivatives (1), (2) of this invention and their salts possess higher antibacterial activities (bactericidal action or antibacterial action) as compared with the hitherto known biguanide derivatives, and some of them also possess antiviral activity. Furthermore, the biguanide derivatives (1), (2) of the invention and their salts possess a broad antibacterial spectrum, and are also excellent in rapidity of action, stability, safety and others, and their activities last for a long time. In addition, they are stimulation-free, odor-free, and soluble also in water. In particular, the compounds in Examples 1 to 4 mentioned below and their salts are particularly excellent in these features, and above all the compound of Example 1 and its salt are excellent in, including the above features, broad antibacterial spectrum, high short-time bactericidal activity, long-lasting drug effect, no problem in color, odor and taste, low toxicity, and the like. The salt of the compound of this invention is excellent in various points, including the above features, such as high solubility, low stimulation and toxicity, high stability, and the like.

Therefore, the biguanide derivatives (1), (2) of this invention and their salts are useful as active ingredient for disinfectants for humans, animals, and medical appliances. The disinfectant of this invention is used in a form of solution, dispersion or suspension by dissolving, dispersing or suspending a specified amount of the biguanide derivatives (1), (2) or their salts in water or organic solvent. Typical examples are eye-drop, nose-drop, gargle, detergent, cleaning agent, and other external application liquids. In these cases, the content of the biguanide derivatives (1), (2) or their salts may be usually about 0.01 to 20 % by weight of the total amount.

Besides, the biguanide derivatives (1), (2) of this invention and their salts may be contained in various cosmetics, such as creams, lotions, powders, colors, makeups, toothpaste, shampoo, soap, depilatories, bleaches, hair-dyes, hair tonics, bath additives, manicure, antiperspiration agent, deodorant, aerosol cosmetics, and baby cosmetics, and the like.

Cosmetics are manufactured by dissolving, dispersing or suspending the specified amount of the biguanide derivatives (1), (2) or their salts, together with other ingredients, in water, other solvent, or various cosmetic bases. The content of the biguanide derivative (1), (2) or their salts in the cosmetic is usually about 0.001 to 1 % by weight of the total amount.

EXAMPLES

Referring to examples, the biguanide derivative of this invention and the disinfectant containing the derivative are explained in detail below. It must be, however, noted that this invention is not limited to these examples.

Example 1

Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-octyl-biguanide hydrochloride

To 20 g of N¹-cyano-N³-octyl-guanidine and 20.2 g of 3,4-dichlorobenzylamine hydrochloride, 200 ml of mesitylene was added, which was heated and refluxed for 1.5 hours. Returning to room temperature after reaction, mesitylene was removed. To the residue, 200 ml of 10% ethanol aqueous solution was added,

which was heated, and stirred for 3 hours at room temperature to be solidified. It was filtered off, and washed with 10% ethanol solution, water and isopropylether in turn, and 28.1 g of rough product was obtained. It was recrystallized in ethyl acetate, and 22.1 g of the captioned compound was obtained. White edged crystals m p. 177-179 °C

Elemental analysis $C_{17}H_{28}Cl_3N_5$			
	C	H	N
Calcd.	49.95	6.90	17.13
Found	49.67	7.18	17.01

Example 2

Preparation of N¹-(4-chlorophenyl)-N⁵-(3,4-dichlorobenzyl)-biguanide hydrochloride

To 20 g of N¹-cyano-N³-(3,4-dichlorobenzyl)-guanidine and 13.5 g of 4-chloroaniline hydrochloride, 200 ml of water was added, which was heated and refluxed for 2 hours. When cooled gradually after reaction, crystals were precipitated. After filtering the crystals, by drying the crystals obtained by heat treatment in 200 ml of ethyl acetate, 21.8 g of the captioned compound was obtained. White edged crystals mp. 238-240 °C

Elemental analysis $C_{15}H_{15}Cl_4N_5$			
	C	H	N
Calcd.	44.25	3.71	17.20
Found	44.19	3.66	17.17

Example 3

Preparation of N¹-(3,4-dichlorophenyl)-N⁵-octylbiguanide hydrochloride

In the same procedure as in Example 1 except that 3,4-dichlorobenzylamine hydrochloride was replaced by the equimolecular amount of 3,4-dichloroaniline hydrochloride, 28.2 g of the captioned compound was obtained. White edged crystals mp. 176-177 °C

Elemental analysis $C_{16}H_{26}Cl_2N_5$			
	C	H	N
Calcd.	48.68	6.64	17.74
Found	48.52	6.50	17.77

Example 4

Preparation of N¹-benzyl-N⁵-dodecylbiguanide hydrochloride

To 20 g of N¹-cyano-N³-benzyl-guanidine and 25.4 g of dodecylamine hydrochloride, 200 ml of mesitylene was added, which was heated and refluxed for 2 hours. After reaction, by the same operation as in Example 1, 23.7 g of the captioned compound was obtained. White edged crystals mp. 135-137 °C

EP 0 507 317 A2

Elemental analysis $C_{21}H_{38}ClN_5$			
	C	H	N
Calcd.	63.69	9.67	17.68
Found	63.47	9.41	17.73

Example 5

Preparation of N¹-benzyl-N⁵-decyl-biguanide hydrochloride

In the same manner as in Example 4 except that dodecylamine hydrochloride was replaced by the equimolecular amount of decylamine hydrochloride, 20.3 g of the captioned compound was obtained.

White edged crystals mp. 133–135 °C

Elemental analysis $C_{19}H_{34}ClN_5$			
	C	H	N
Calcd.	62.02	9.31	19.03
Found	62.29	9.57	18.91

Example 6

Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-(3-trifluoromethylphenyl)-biguanide hydrochloride

In the same manner as in Example 2 except that 4-chloroaniline hydrochloride was replaced by the equimolecular amount of 3-trifluoromethylaniline hydrochloride, 23.2 g of the captioned compound was obtained.

White edged crystals mp. 185–188 °C

Elemental analysis $C_{16}H_{15}Cl_2F_3N_5$			
	C	H	N
Calcd.	43.61	3.43	15.89
Found	43.57	3.24	16.10

Example 7

Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-(4-bromophenyl)-biguanide hydrochloride

In the same manner as in Example 2 except that 4-chloroaniline hydrochloride was replaced by the equimolecular amount of 4-bromoaniline hydrochloride, 21.6 g of the captioned compound was obtained.

White edged crystals mp. 223–225 °C

Elemental analysis $C_{15}H_{15}BrCl_2N_5$			
	C	H	N
Calcd.	39.90	3.35	15.51
Found	40.34	3.16	15.82

Example 8

Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-(4-iodophenyl)-biguanide hydrochloride

In the same manner as in Example 2 except that 4-chloroaniline hydrochloride was replaced by the equimolecular amount of 4-iodoaniline hydrochloride, 22.6 g of the captioned compound was obtained.

5 White edged crystals mp. 215~217° C

Elemental analysis C ₁₅ H ₁₅ Cl ₂ IN ₅			
	C	H	N
Calcd.	36.14	3.03	14.05
Found	36.25	2.80	14.21

15 Example 9

Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-(2,4-dichlorophenyl)-biguanide hydrochloride

20 In the same manner as in Example 2 except that 4-chloroaniline hydrochloride was replaced by the equimolecular amount of 2,4-dichloroaniline hydrochloride, 20.4 g of the captioned compound was obtained. White edged crystals mp. 217~220° C

Elemental analysis C ₁₅ H ₁₅ Cl ₄ N ₅			
	C	H	N
Calcd.	40.80	3.20	15.86
Found	41.25	2.93	15.79

30 Example 10

Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-(3,4-dichlorophenyl)-biguanide hydrochloride

35 In the same manner as in Example 2 except that 4-chloroaniline hydrochloride was replaced by the equimolecular amount of 3,4-dichloroaniline hydrochloride, 21.7 g of the captioned compound was obtained. White edged crystals mp. 205~208° C

Elemental analysis C ₁₅ H ₁₄ Cl ₄ N ₅			
	C	H	N
Calcd.	40.80	3.20	15.86
Found	41.03	2.94	15.94

45 Example 11

Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-(2,3,4-trichlorophenyl)-biguanide hydrochloride

50 In the same manner as in Example 2 except that 4-chloroaniline hydrochloride was replaced by the equimolecular amount of 2,3,4-trichloroaniline hydrochloride, 23.2 g of the captioned compound was obtained.

55 White edged crystals mp. 227~228° C

Elemental analysis $C_{15}H_{13}Cl_6N_5$			
	C	H	N
Calcd.	37.85	2.75	14.71
Found	38.09	2.55	14.81

Example 12

Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-(3,4-dimethylphenyl)-biguanide hydrochloride

In the same manner as in Example 2 except that 4-chloroaniline hydrochloride was replaced by the equimolecular amount of 3,4-dimethylaniline hydrochloride, 20.3 g of the captioned compound was obtained.

White edged crystals mp. 190~191 °C

Elemental analysis $C_{17}H_{20}Cl_3N_5$			
	C	H	N
Calcd.	50.95	5.03	17.48
Found	50.66	4.78	17.56

Example 13

Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-(3,4-methylenedioxyphenyl)-biguanide hydrochloride

In the same manner as in Example 2 except that 4-chloroaniline hydrochloride was replaced by the equimolecular amount of 3,4-methylenedioxyaniline hydrochloride, 25.7 g of the captioned compound was obtained.

White edged crystals mp. 204~206 °C

Elemental analysis $C_{16}H_{16}Cl_2N_5O_2$			
	C	H	N
Calcd.	46.12	3.87	16.81
Found	45.83	3.69	16.97

Example 14

Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-(4-t-butylphenyl)-biguanide hydrochloride

In the same manner as in Example 2 except that 4-chloroaniline hydrochloride was replaced by the equimolecular amount of 4-t-butylaniline hydrochloride, 22.5 g of the captioned compound was obtained.

White edged crystals mp. 228~230 °C

Elemental analysis $C_{19}H_{24}Cl_2N_5$			
	C	H	N
Calcd.	53.22	5.64	16.33
Found	52.93	5.38	16.16

Example 15

Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-(4-ethyl-thiophenyl)-biguanide hydrochloride

- 5 In the same manner as in Example 2 except that 4-chloroaniline hydrochloride was replaced by the equimolecular amount of 4-ethylthioaniline hydrochloride, 20.7 g of the captioned compound was obtained. White edged crystals mp. 199~200 °C

10

Elemental analysis C ₁₇ H ₂₀ Cl ₂ N ₅ S			
	C	H	N
Calcd.	47.18	4.66	16.18
Found	47.12	4.46	16.04

15

Example 16

Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-(n-decyl)-biguanide hydrochloride

20

- In the same manner as in Example 1 except that N¹-cyano-N³-octyl-guanidine was replaced by the equimolecular amount of N¹-cyano-N³-(n-decyl)-guanidine, 24.5 g of the captioned compound was obtained. White edged crystals mp. 135~137 °C

25

Elemental analysis C ₁₉ H ₃₂ Cl ₂ N ₅			
	C	H	N
Calcd.	52.24	7.38	16.03
Found	52.08	7.12	16.31

30

Example 17

Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-(1,1,3,3-tetramethylbutyl)-biguanide hydrochloride

35

- In the same manner as in Example 1 except that N¹-cyano-N³-octyl-guanidine was replaced by the equimolecular amount of N¹-cyano-N³-(1,1,3,3-tetrabutyl)-guanidine, 23.1 g of the captioned compound was obtained.

- 40 White edged crystals mp. 196~198 °C

45

Elemental analysis C ₁₇ H ₂₈ Cl ₂ N ₅			
	C	H	N
Calcd.	49.95	6.90	17.13
Found	49.78	6.81	17.01

50 Example 18

Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-hexyl-biguanide hydrochloride

- 55 In the same manner as in Example 1 except that N¹-cyano-N³-octyl-guanidine was replaced by the equimolecular amount of N¹-cyano-N³-hexyl guanidine, 19.7 g of the captioned compound was obtained. White edged crystals mp. 155~157 °C

Elemental analysis $C_{15}H_{24}Cl_3N_5$			
	C	H	N
Calcd.	47.32	6.35	18.39
Found	47.21	6.17	18.50

Example 19

N¹-(3,4-dichlorobenzyl)-N⁵-(2-ethoxyethyl)-biguanide hydrochloride

In the same manner as in Example 1 except that N¹-cyano-N³-octyl-guanidine was replaced by the equimolecular amount of N¹-cyano-N³-(2-ethoxyethyl)-guanidine, 23.1 g of the captioned compound was obtained.

White edged crystals mp. 170~173° C

Elemental analysis $C_{13}H_{20}Cl_3N_5O$			
	C	H	N
Calcd.	42.35	5.47	18.99
Found	42.09	5.18	18.73

Example 20

N¹-(3,4-dichlorobenzyl)-N⁵-[2-(2-hydroxyethoxy)ethyl]-biguanide hydrochloride

In the same manner as in Example 1 except that N¹-cyano-N³-octyl-guanidine was replaced by the equimolecular amount of N¹-cyano-N³-[2-(2-hydroxyethoxy)ethyl]-guanidine, 21.5 g of the captioned compound was obtained.

White edged crystals mp. 140~142° C

Elemental analysis $C_{13}H_{20}Cl_3N_5O_2$			
	C	H	N
Calcd.	40.59	5.24	18.20
Found	40.30	5.01	18.03

Example 21

Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-(3-diethylaminopropyl)-biguanide

A rough product was obtained in the same manner as in Example 1 except that N¹-cyano-N³-octyl-guanidine was replaced by the equimolecular amount of N¹-cyano-N³-(3-diethylaminopropyl)-guanidine. The product was dissolved in 100 ml of methanol, and an equimolecular amount of 1N sodium hydroxide was added, which was concentrated at reduced pressure, and the residue was recrystallized in isopropanol, and 16.7 of the captioned compound was obtained.

White edged crystals mp. 156~158° C

Elemental analysis $C_{16}H_{26}Cl_2N_6$			
	C	H	N
Calcd.	51.48	7.02	22.51
Found	51.21	6.83	22.76

In the following Examples 22 to 26, using proper starting materials, the individual compounds were prepared in the same manner as in Example 1.

Example 22

Preparation of N¹-(3,4-Dichlorobenzyl)-N⁵-(3-(2-ethylhexyloxy)-propyl)-biguanide hydrochloride

White edged crystals mp. 200~203 ° C

Elemental analysis $C_{20}H_{34}Cl_2N_5O$			
	C	H	N
Calcd.	51.45	7.34	15.00
Found	51.12	7.62	14.82

Example 23

Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-[(3-isopropoxy)propyl]-biguanide hydrochloride

White edged crystals mp. 158~160 ° C

Elemental analysis $C_{15}H_{24}Cl_2N_5O$			
	C	H	N
Calcd.	45.41	6.10	17.65
Found	45.17	6.02	17.53

Example 24

Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-[(2-diethylamino)-ethyl]-biguanide

White edged crystals mp. 137~140 ° C

Elemental analysis $C_{15}H_{24}Cl_2N_6$			
	C	H	N
Calcd.	50.14	6.73	23.39
Found	50.01	6.48	23.15

Example 25

Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-[(3-butyl)-propyl]-biguanide hydrochloride

White edged crystals mp. 181~184 ° C

EP 0 507 317 A2

Elemental analysis C ₁₆ H ₂₆ Cl ₃ N ₅ O			
	C	H	N
Calcd.	46.78	6.38	17.05
Found	46.49	6.27	17.21

Example 26

Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-[3-(di-n-butylamino)propyl]-biguanide dihydrochloride

White edged crystals mp. 122~124 ° C

Elemental analysis C ₂₀ H ₃₆ Cl ₄ N ₆			
	C	H	N
Calcd.	47.82	7.22	16.73
Found	47.72	7.01	16.65

Example 27

Preparation of N¹-(4-chlorophenyl)-N⁵-cyclohexylmethyl-biguanide hydrochloride

In the same manner as in Example 2 except that N¹-cyano-N³-(3,4-dichlorobenzyl)-guanidine was replaced by the equimolecular amount of N¹-cyano-N³-cyclohexylmethyl-guanidine, 20.4 g of the captioned compound was obtained.

White edged crystals mp. 240~241 ° C

Elemental analysis C ₁₅ H ₂₃ Cl ₂ N ₅			
	C	H	N
Calcd.	52.33	6.73	20.34
Found	52.24	6.61	20.48

Example 28

Preparation of N¹-(4-chlorophenyl)-N⁵-(3-trifluoromethylphenyl)-biguanide hydrochloride

In the same manner as in Example 2 except that N¹-cyano-N³-(3,4-dichlorobenzyl)-guanidine was replaced by the equimolecular amount of N¹-cyano-N³-(3-trifluoromethylphenyl)-guanidine, 22.7 g of the captioned compound was obtained.

White edged crystals mp. 213~215 ° C

Elemental analysis C ₁₅ H ₁₄ Cl ₂ F ₃ N ₅			
	C	H	N
Calcd.	45.93	3.60	17.86
Found	46.03	3.55	18.04

Example 29

Preparation of N¹-(4-chlorophenyl)-N⁵-(4-ethylthiophenyl)-biguanide hydrochloride

In the same manner as in Example 2 except that N¹-cyano-N³-(3,4-dichlorobenzyl)-guanidine was replaced by the equimolecular amount of N¹-cyano-N³-(4-ethylthiophenyl)-guanidine, 20.76 g of the captioned compound was obtained.

White edged crystals mp. 243~245 ° C

Elemental analysis C ₁₆ H ₁₉ Cl ₂ N ₅ S			
	C	H	N
Calcd.	50.00	4.98	18.22
Found	50.04	4.82	18.10

Example 30

Preparation of N¹-(4-chlorophenyl)-N⁵-(4-chlorobenzyl)-biguanide hydrochloride

In the same manner as in Example 2 except that N¹-cyano-N³-(3,4-dichlorobenzyl)-guanidine was replaced by the equimolecular amount of N¹-cyano-N³-(4-chlorobenzyl)-guanidine, 21.6 g of the captioned compound was obtained.

White edged crystals mp. 225~227 ° C

Elemental analysis C ₁₅ H ₁₆ Cl ₂ N ₅			
	C	H	N
Calcd.	48.34	4.33	18.79
Found	48.32	4.19	18.86

Example 31

Preparation of N¹-(4-chlorophenyl)-N⁵-(2,4-dichlorobenzyl)-biguanide hydrochloride

In the same manner as in Example 2 except that N¹-cyano-N³-(3,4-dichlorobenzyl)-guanidine was replaced by the equimolecular amount of N¹-cyano-N³-(2,4-dichlorobenzyl)-guanidine, 20.1 g of the captioned compound was obtained.

White edged crystals mp. 234~236 ° C

Elemental analysis C ₁₅ H ₁₅ Cl ₄ N ₅			
	C	H	N
Calcd.	44.25	3.71	17.20
Found	44.42	3.50	16.95

Example 32

Preparation of N¹-(4-chlorophenyl)-N⁵-(4-acetylaminophenyl)-biguanide hydrochloride

In the same manner as in Example 2 except that N¹-cyano-N³-(3,4-dichlorobenzyl)-guanidine was replaced by the equimolecular amount of N¹-cyano-N³-(4-acetylaminophenyl)-guanidine, 18.6 g of the captioned compound was obtained.

White edged crystals mp. 253~255 ° C

Elemental analysis $C_{16}H_{18}Cl_2N_6O$			
	C	H	N
Calcd.	50.40	4.76	22.04
Found	50.63	4.51	21.92

Example 33

Preparation of N¹-(4-chlorophenyl)-N⁵-(3,4-methylenedioxyphenyl)-biguanide hydrochloride

In the same manner as in Example 2 except that N¹-cyano-N³-(3,4-dichlorobenzyl)-guanidine was replaced by the equimolecular amount of N¹-cyano-N³-(3,4-methylenedioxyphenyl)-guanidine, 19.7 g of the captioned compound was obtained.

White edged crystals mp. 237~239 °C

Elemental analysis $C_{15}H_{15}Cl_2N_5O_2$			
	C	H	N
Calcd.	48.93	4.11	8.69
Found	48.76	3.95	8.47

Example 34

Preparation of N¹-(4-chlorophenyl)-N⁵-(3,4-methylenedioxybenzyl)-biguanide hydrochloride

In the same manner as in Example 2 except that N¹-cyano-N³-(3,4-dichlorobenzyl)-guanidine was replaced by the equimolecular amount of N¹-cyano-N³-(3,6-methylenedioxybenzyl)-guanidine, 22.4 g of the captioned compound was obtained.

White edged crystals mp. 231~233 °C

Elemental analysis $C_{16}H_{17}Cl_2N_5O_2$			
	C	H	N
Calcd.	50.27	4.48	18.32
Found	50.44	4.29	18.31

Example 35

Preparation of N¹-(4-chlorobenzyl)-N⁵-octyl-biguanide hydrochloride

In the same manner as in Example 1 except that 3,4-dichlorobenzylamine hydrochloride was replaced by the equimolecular amount of 2,4-chlorobenzylamine hydrochloride, 21.3 g of the captioned compound was obtained.

White edged crystals mp. 156~158 °C

Elemental analysis $C_{17}H_{29}Cl_2N_5$			
	C	H	N
Calcd.	54.54	7.81	18.71
Found	54.76	7.68	18.69

Example 36

Preparation of N¹-(4-chlorobenzyl)-N⁵-(4-chlorobenzyl)-biguanide hydrochloride

- 5 Using 20.85 g of N¹-cyano-N⁵-(4-chlorobenzyl)-guanidine and 17.8 g of 4-chlorobenzylamine hydrochloride, 21.4 g of the captioned compound was obtained by operating the same manner as in Example 1. White edged crystals mp. 200~202 °C

10

Elemental analysis C ₁₆ H ₁₈ Cl ₂ N ₅			
	C	H	N
Calcd.	49.69	4.69	18.11
Found	49.62	4.61	18.37

15

Example 37

Preparation of N¹-(4-chlorobenzyl)-N⁵-decyl-biguanide hydrochloride

20

In the same manner as in Example 36 except that 4-chlorobenzylamine hydrochloride was replaced by the equimolecular amount of decylamine hydrochloride, 24.3 g of the captioned compound was obtained. White edged crystals mp. 161~163 °C

25

Elemental analysis C ₁₉ H ₃₃ Cl ₂ N ₅			
	C	H	N
Calcd.	56.71	8.27	17.40
Found	56.60	8.12	17.63

30

Example 38

Preparation of N¹-(4-chlorobenzyl)-N⁵-dodecyl-biguanide hydrochloride

35

In the same manner as in Example 36 except that 4-chlorobenzylamine hydrochloride was replaced by the equimolecular amount of dodecylamine hydrochloride, 22.3 g of the captioned compound was obtained. White edged crystals mp. 158~160 °C

40

Elemental analysis C ₂₁ H ₃₇ Cl ₂ N ₅			
	C	H	N
Calcd.	58.59	8.66	16.27
Found	58.41	8.53	16.55

45

Example 39

50

Preparation of N¹-(4-chlorobenzyl)-N⁵-isobutyl-biguanide hydrochloride

- In the same manner as in Example 36 except that 4-chlorobenzylamine hydrochloride was replaced by the equimolecular amount of isobutylamine hydrochloride, 18.7 g of the captioned compound was obtained.
- 55 White edged crystals mp. 219~221 °C

EP 0 507 317 A2

Elemental analysis C ₁₃ H ₂₁ Cl ₂ N ₅			
	C	H	N
Calcd.	49.06	6.65	22.01
Found	48.87	6.52	22.23

Example 40

Preparation of N¹-(4-chlorobenzyl)-N⁵-(3,4-dichlorophenyl)-biguanide hydrochloride

In the same manner as in Example 36 except that 4-chlorobenzylamine hydrochloride was replaced by the equimolecular amount of 3,4-dichloraniline hydrochloride, 21.7 g of the captioned compound was obtained.

White edged crystals mp. 183~185 °C

Elemental analysis C ₁₅ H ₁₅ Cl ₄ N ₅			
	C	H	N
Calcd.	44.25	3.71	17.20
Found	44.37	3.66	17.42

Example 41

Preparation of N¹-(3,4-dichlorophenyl)-N⁵-(3,4-methylenedioxybenzyl)-biguanide hydrochloride

Using 21.5 g of N¹-cyano-N³-(3,4-dichlorophenyl)-guanidine and 18.8 g of 3,4-methylenedioxybenzylamine hydrochloride, 27.3 g of the captioned compound was obtained by operating the same manner as in Example 1.

White edged crystals mp. 175~177 °C

Elemental analysis C ₁₆ H ₁₆ Cl ₂ N ₅ O ₂			
	C	H	N
Calcd.	46.12	3.87	16.81
Found	46.01	3.76	16.97

Example 42

Preparation of N¹-(3,4-dichlorophenyl)-N⁵-hexyl-biguanide hydrochloride

In the same manner as in Example 41 except that 3,4-methylenedioxybenzylamine hydrochloride was replaced by the equimolecular amount of hexylamine hydrochloride, 22.3 g of the captioned compound was obtained.

White edged crystals mp. 154~156 °C

Elemental analysis C ₁₄ H ₂₂ Cl ₂ N ₅			
	C	H	N
Calcd.	45.85	6.05	19.10
Found	45.66	6.17	19.03

Example 43

Preparation of N¹-(3,4-dichlorophenyl)-N⁵-decyl-biguanide hydrochloride

5 In the same manner as in Example 41 except that 3,4-methylenedioxybenzylamine hydrochloride was replaced by the equimolecular amount of decylamine hydrochloride, 24.2 g of the captioned compound was obtained.

White edged crystals mp. 131~133 °C

10

Elemental analysis C ₁₈ H ₃₀ Cl ₂ N ₅			
	C	H	N
Calcd.	51.13	7.15	6.56
Found	51.02	7.32	16.37

15

Example 44

20 Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-octyl-biguanide lactate

Dissolving 20 g of N¹-(3,4-dichlorobenzyl)-N⁵-octyl-biguanide hydrochloride in 200 ml of methanol, 24.5 ml of 5N NaOH was added, and the solvent was distilled away. The residual white solid was washed with water, suspended in 200 ml of water off, and 13 g of lactic acid was added, and dissolved by heating.
 25 Chilled in ice, precipitating crystals were filtered, and 12.14 g of the captioned compound was obtained.
 White edged crystals m p . 45~46 °C

Example 45

30 Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-octyl-biguanide glycolate

The captioned compound was obtained in the same manner as in Example 44 except that lactic acid was replaced by glycolic acid.
 White edged crystals mp. 109~110 °C

35

Example 46

Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-octyl-biguanide monomethanesulfonate

40 The captioned compound was obtained in the same manner as in Example 44 except that lactic acid was replaced by methanesulfonic acid.
 White edged crystals mp. 174~176 °C

Example 47

45

Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-octyl-biguanide hydrobromide

The captioned compound was obtained in the same manner as in Example 44 except that lactic acid was replaced by 47% hydrobromic acid.
 50 White edged crystals mp. 120~121 °C

Example 48

55 Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-octyl-biguanide phosphate

The captioned compound was obtained in the same manner as in Example 44 except that lactic acid was replaced by phosphoric acid.
 White edged crystals mp. 96~98 °C

Example 49

Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-octyl- biguanide dimethanesulfonate

5 Dissolving 25 g of N¹-(3,4-dichlorobenzyl)-N⁵-octyl-biguanide hydrochloride in 250 ml of methanol, 30.6 ml of 5N NaOH was added, and the solvent was distilled away. The residual white solid was washed with water, and dissolved in 500 ml of acetone, and 14.7 g of methanesulfonic acid was added. The precipitating crystals were filtered off, and by recrystallizing from ethanol-ether, 25.8 g of the captioned compound was obtained.

10 White edged crystals mp. 171~172 °C

Elemental analysis C ₁₇ H ₂₇ N ₅ Cl ₂ · 2CH ₃ SO ₃ H			
	C	H	N
Calcd.	40.42	6.25	12.41
Found	40.40	6.55	12.31

20 Reference Example

Preparation of N¹-(3,4-dichlorobenzyl)-N³-cyanoguanidine

25 Putting 63.7 g (300 mM) of 3,4-dichlorobenzylamine hydrochloride and 30.9 g (1.1 eq.) of sodium dicyanamide in 950 ml (15 V) of acetonitrile, the mixture was heated and refluxed for 3.5 hours. Distilling away the solvent, 600 ml of water was added to the residue (white solid) to disperse and wash. Obtained crystals were filtered off, and dispersed and washed in 600 ml of dichloromethane. Since residue of amine was recognized, the crystals were dispersed and washed again in 600 ml of water and 500 ml of dichloromethane. Crystals were filtered off, and dried overnight at 60 °C, and 57.7 g (78.9%) of the

30 captioned compound was obtained.

White crystals mp. 173 °C

Elemental analysis C ₉ H ₈ Cl ₂ N ₄			
	C	H	N
Calcd.	44.47	3.32	23.05
Found	44.32	3.08	23.30

40 Example 50

Preparation of N¹-(3,4-dichlorobenzyl)-N⁵-octyl-biguanide dihydrochloride

45 Suspending 10.0 g (41.1 mM) of N¹-(3,4-dichlorobenzyl)-N³-cyanoguanidine and 5.31 g (1.0 eq.) of n-octylamine in 100 ml of xylene, 3.5 ml (1.1 eq.) of concentrated hydrochloric acid was added and stirred for several minutes. The reaction was continued for 8 hours in reflux by using Dean-Stark trap. Distilling away the solvent, and by ethanol coboiling, xylene was removed. The residue was dissolved in 40 ml of 70% acetonitrile, and 6.8 ml (2.0 eq.) of concentrated hydrochloric acid was added. Further adding 90 ml of 70%

50 acetonitrile, the mixture was recrystallized. Obtained crystals were filtered off, dried overnight at 60 °C, and 14.3 g (78.2%) of white crystalline product was obtained. The obtained rough crystals were recrystallized twice with 70% acetonitrile (6 to 7 V), and 7.16 g (50.1%) of the captioned compound was obtained.

mp. 176~178 °C

55

Elemental analysis $C_{17}H_{29}Cl_4N_5$			
	C	H	N
Calcd.	45.86	6.56	15.73
Found	45.54	6.66	15.79

5

Meanwhile, when the produced material (dihydrochloride) was recrystallized in a solvent of high water content (such as CH_3CN , ethanol), it was converted into the same monohydrochloride as obtained in Example 1.

10

Prescription 1 of disinfectant

15	Monobiguanide derivative of Example 1	5 g
	Nonionic surface active agent	3.75 g
	(Polyoxyethylene phenylether)	
20	<u>Distilled water for injection</u>	<u>proper</u>
	Whole volume	100 ml

Prescription 2 of disinfectant

25	Monobiguanide derivative of Example 1	0.5 g
	Nonionic surface active agent	0.375 g
30	(Polyoxyethylene phenylether)	
35	Ethanol	83 ml
	<u>Distilled water for injection</u>	<u>proper</u>
40	Whole volume	100 ml

[Antibacterial activity test]

45 In order to test the antibacterial action of the compounds obtained in the examples in various organisms, the minimum inhibitory concentration (MIC) was determined according to the standard method of Japan Society of Chemotherapy. The culture medium for sensitivity disc was used (Chemotherapy, vol. 29, No. 1, 76-79, 1981). As the control drug, gluconic chlorohexydine (tradename "Hibiten" commercially available from Sumitomo Pharmaceutical) was used and tested similarly. The results are shown in Table 1.

50 The inoculum size was adjusted to 1×10^6 cells/ml (0.07 to 0.16 at O.D. 600 nm).

55

Table 1 (1/7)

Strain	MIC $\mu\text{g/ml}$ (10^6 cells/ml)						
	Examples						
	1	2	3	4	5	6	7
<i>S. aureus</i> FDA 209P	1.56	1.56	3.13	1.56	0.78	3.13	1.56
<i>S. aureus</i> MRSA 57	1.56	3.13	3.13	1.56	1.56	3.13	1.56
<i>E. coli</i> NIHJ JC-2	100	12.5	6.25	100	50	12.5	6.25
<i>K. pneumoniae</i> NCTC 9632	—	—	—	—	—	—	—
<i>S. marcescens</i> IFO 12648	100	25	100	100	50	25	25
<i>P. aeruginosa</i> ATCC 10145	100	25	100	100		100	50
<i>A. calcoaceticus</i> Ac-54	25	12.5	12.5	100	25	12.5	6.25

Table 1 (2/7)

Strain	MIC $\mu\text{g}/\text{ml}$ (10^6 cells/ml)												
	Examples												
	8	9	10	11	12	13	14						
<i>S. aureus</i> FDA 209P	1.56	1.56	0.78	1.56	1.56	12.5	1.56						
<i>S. aureus</i> MRSA 57	1.56	1.56	0.39	0.78	1.56	12.5	1.56						
<i>E. coli</i> NIHJ JC-2	12.5	25	6.25	6.25	12.5	50	100						
<i>K. pneumoniae</i> NCTC 9632	—	—	—	—	—	—	—						
<i>S. marcescens</i> IFO 12648	25	25	6.25	12.5	25	50	50						
<i>P. aeruginosa</i> ATCC 10145	100	100	25	50	50	100	100						
<i>A. calcoaceticus</i> Ac-54	12.5	6.25	3.13	6.25	12.5	100	6.25						

Table 1 (3/7)

Strain	MIC $\mu\text{g}/\text{ml}$ (10^6 cells/ml)									
	Examples									
	15	16	17	18	19	20	21			
<i>S. aureus</i> FDA 209P	3.13	1.56	1.56	0.78	6.25	6.25	12.5			
<i>S. aureus</i> MRSA 57	3.13	1.56	1.56	0.78	6.25	6.25	12.5			
<i>E. coli</i> NIHJ JC-2	25	100	100	12.5	25	25	100			
<i>K. pneumoniae</i> NCTC 9632	—	—	—	—	—	—	—			
<i>S. marcescens</i> IFO 12648	100	100	100	12.5	50	50	100			
<i>P. aeruginosa</i> ATCC 10145	100	100	100	100	100	100	100			
<i>A. calcoaceticus</i> Ac-54	12.5	100	12.5	6.25	50	50	100			

Table 1 (4/7)

Strain	MIC $\mu\text{g}/\text{ml}$ (10^6 cells/ml)					
	Examples					
	22	23	24	25	26	27
<i>S. aureus</i> FDA 209P	1. 56	3. 13	6. 25	1. 56	6. 25	6. 25
<i>S. aureus</i> MRSA 57	1. 56	3. 13	6. 25	1. 56	6. 25	6. 25
<i>E. coli</i> NIHJ JC-2	100	50	100	25	50	25
<i>K. pneumoniae</i> NCTC 9632	—	—	—	—	25	—
<i>S. marcescens</i> IFO 12648	100	50	100	25	100	50
<i>P. aeruginosa</i> ATCC 10145	100	100	100	100	100	100
<i>A. calcoaceticus</i> Ac-54	100	50	100	25	100	12. 5

Table 1 (5/7)

Strain	MIC $\mu\text{g}/\text{ml}$ (10^6 cells/ml)						
	Examples						
	28	29	30	31	32	33	34
<i>S. aureus</i> FDA 209P	6.25	6.25	3.13	1.56	25	50	25
<i>S. aureus</i> MRSA 57	6.25	6.25	0.78	0.78	12.5	50	25
<i>E. coli</i> NIHJ JC-2	25	12.5	12.5	12.5	100	100	100
<i>K. pneumoniae</i> NCTC 9632	—	—	—	—	—	—	—
<i>S. marcescens</i> IFO 12648	50	25	12.5	12.5	100	100	100
<i>P. aeruginosa</i> ATCC 10145	100	100	25	25	100	100	100
<i>A. calcoaceticus</i> Ac-54	12.5	12.5	6.25	6.25	100	100	100

Table 1 (6/7)

Strain	MIC $\mu\text{g}/\text{ml}$ (10^6 cells/ml)									
	Examples									
	35	36	37	38	39	40	41			
<i>S. aureus</i> FDA 209P	3.13	3.13	1.56	6.25	12.5	1.56	12.5			
<i>S. aureus</i> MRSA 57	3.13	3.13	1.56	6.25	12.5	1.56	6.25			
<i>E. coli</i> NIHJ JC-2	100	25	100	100	50	12.5	25			
<i>K. pneumoniae</i> NCTC 9632	—	—	—	—	50	—	—			
<i>S. marcescens</i> IFO 12648	100	25	100	100	100	12.5	25			
<i>P. aeruginosa</i> ATCC 10145	100	100	100	100	100	100	100			
<i>A. calcoaceticus</i> Ac-54	25	25	100	100	50	6.25	25			

Table 1 (7/7)

Strain	MIC $\mu\text{g}/\text{ml}$ (10^6 cells/ml)							
	Examples							Control agent
	42	43	44	45	46	49		
<i>S. aureus</i> FDA 209P	3.13	6.25	1.56	1.56	1.56	1.56	1.56	1.56
<i>S. aureus</i> MRSA 57	3.13	6.25	1.56	1.56	3.13	3.13	1.56	1.56
<i>E. coli</i> NIHJ JC-2	12.5	100	100	100	100	100	1.56	1.56
<i>K. pneumoniae</i> NCTC 9632	—	50	—	—	—	—	—	—
<i>S. marcescens</i> IFO 12648	6.25	100	100	100	100	100	25	25
<i>P. aeruginosa</i> ATCC 10145	6.25	100	100	100	100	100	100	100
<i>A. calcoaceticus</i> Ac-54	12.5	100	25	25	25	50	12.5	12.5

[Bactericidal activity test]

As to the compounds obtained in Examples 1 and 2, the bactericidal activity at room temperature (25 °C) was measured by treating in a short time by reference to the phenol coefficient measurement method.

(1) Culture medium

For prior culture of test organism, Mueller Hinton Broth (DIFCD) was used. As growth culture medium of surviving cells in the test solution after bactericidal treatment, SCDLP culture medium inactivated by disinfectant was used ("Daigo" commercially available from Nippon Seiyaku).

(2) Test method

The compound obtained in each examples was dissolved in distilled water, and adjusted to a double concentration of the test concentration, and after aseptic filtering, the solution was diluted in steps of 1/2 in a range of specified test concentration, and 50 μ l was dispensed in 96-well microplate. On the other hand, the test organism undergoing the prior culture for 16 hours twice was adjusted to 10^8 cells/ml with sterilized distilled water in order to eliminate the effects of culture medium (0.3 at O.D. 660 nm), and diluted to 1/100 with 5 ml or 10 ml of sterilized distilled water to obtain 10^6 cells/ml, and into 50 μ l of the test solution in the microplate, an equivalent 50 μ l was injected and mixed. After the lapse of 1 minute and 3 minutes, 5 μ l of the test solution was taken out from the microplate, and was suspended in 150 μ l of inactivated culture medium, and was cultivated for 16 to 18 hours at 37 °C. The survival of organisms in the test solution was judged by the turbidity, and the minimum bactericidal concentration in lapse of 1 minute and 3 minutes was determined. The result is shown in Table 2.

Table 2

Strain	Minimum Bactericidal Concentration $\mu\text{g}/\text{ml}$ (10^6 cells/ml)					
	Example 1		Example 2		Control agent	
	1 min.	3 min.	1 min.	3 min.	1 min.	3 min.
<i>S. aureus</i> FDA 209P	12.5	3.13	500	100	>1000	25
<i>S. aureus</i> MRSA 57	6.25	≤ 1.56	200	25	>1000	50
<i>E. coli</i> NIHJ JC-2	3.13	≤ 1.56	25	6.25	12.5	6.25
<i>K. pneumoniae</i> NCTC 9632	3.13	3.13	12.5	12.5	3.13	≤ 1.56
<i>S. marcescens</i> IFO 12648	6.25	≤ 1.56	25	6.25	12.5	6.25
<i>A. calcoaceticus</i> Ac-54	3.13	3.13	12.5	6.25	6.25	6.25
<i>P. aeruginosa</i> ATCC 10145	3.13	3.13	12.5	6.25	12.5	6.25
<i>P. cepacia</i> 2315-C0010	200	50	200	50	>1000	>1000

Thus, the monobiguanide derivative (1) of the invention or its salt possesses a high bactericidal action and antibacterial action, and hence it may be preferably used as the disinfectant for the human or animal skin or the like.

Next, the examples of bis-biguanide derivative (2) of this invention are explained.

Example 51

Preparation of 1,1'-[1,3-cyclohexanebis(methylene)]-bis[5-(4-chlorophenyl)biguanide]dihydrochloride

Refluxing 33 g (0.15 mols) of 1,3-di(aminomethyl) cyclohexane dihydrochloride and 26 g (0.3 mols) of dicyanamide sodium in butanol for 12 hours at temperature of 140°C , 51.3 g of 1,1'-[1,3-cyclohexanebis-

(methylene)]-bis(3-cyanoguanidine) was obtained.

Adding 5.52 g (0.02 mols) of this reaction product and 6.56 g (0.04 mols) of p-chloroaniline hydrochloride into β -oxyethylether (tradename: Cellosolve), by refluxing for 3 hours at 140 °C to react, 4.3 g of product was obtained. This product was recrystallized and refined in water and ethanol mixed solvent, and the captioned compound was obtained.

White edged crystals mp. 245~248 °C

N M R (CDCl₃ + DMSO-d₆) δ :

1.1 ~ 1.9 (m, 8H), 2.8~ 3.5 (m, 6H), 6.95 (bs), 7.37 (dd, 4H, J = 9 Hz) 7.90 (bs)

Example 52

Preparation of 1,1'-[1,3-cyclohexanebis(methylene)]-bis[5-(4-iodophenyl)biguanide]dihydrochloride

The captioned compound was obtained in the same manner as in Example 51 except that p-chloroaniline hydrochloride was replaced by the equimolecular amount of p-iodoaniline hydrochloride (Yield 5.1 g).

White edged crystals mp. 259~261 °C

Example 53

Preparation of 1,1'-[1,3-cyclohexanebis(methylene)]-bis[5-(4-bromophenyl)biguanide]dihydrochloride

The captioned compound was obtained in the same manner as in Example 51 except that p-chloroaniline hydrochloride was replaced by the equimolecular amount of p-bromoaniline hydrochloride (Yield 4.7 g).

White edged crystals mp. 265~267 °C

Example 54

Preparation of 1,1'-[1,3-cyclohexanebis(methylene)]-[5-(2,4-dichlorophenyl)biguanide]dihydrochloride

The captioned compound was obtained in the same manner as in Example 51 except that p-chloroaniline hydrochloride was replaced by the equimolecular amount of 2,4-dichloroaniline hydrochloride (Yield 5.7 g).

White edged crystals mp. 261~265 °C

Example 55

Preparation of 1,1'-[1,3-cyclohexanebis(methylene)]-[5-(2,4-difluorophenyl)biguanide]dihydrochloride

The captioned compound was obtained in the same manner as in Example 51 except that p-chloroaniline hydrochloride was replaced by the equimolecular amount of 2,4-difluoroaniline hydrochloride (Yield 4.1 g).

White edged crystals mp. 212~215 °C

Example 56

Preparation of 1,1'-[1,3-cyclohexanebis(methylene)]-[5-(4-t-butylphenyl)biguanide]dihydrochloride

The captioned compound was obtained in the same manner as in Example 51 except that p-chloroaniline hydrochloride was replaced by the equimolecular amount of p-t-butylaniline hydrochloride (Yield 3.6 g).

White edged crystals mp. 194~197 °C

Example 57

Preparation of 1,1'-[1,3-cyclohexanebis(methylene)]-[5-(3,4-dimethylphenyl)biguanide]dihydrochloride

The captioned compound was obtained in the same manner as in Example 51 except that p-chloroaniline hydrochloride was replaced by the equimolecular amount of 3,4-dimethylaniline hydrochloride (Yield 3.8 g).

Pale yellow edged crystals mp. 162~166 °C

5

Example 58

Preparation of 1,1'-[1,3-cyclohexanebis(methylene)]-[5-(4-ethylthiophenyl)biguanide]dihydrochloride

10 The captioned compound was obtained in the same manner as in Example 51 except that p-chloroaniline hydrochloride was replaced by the equimolecular amount of p-ethylthioaniline hydrochloride (Yield 5.4 g).

mp. 110~113 °C

15 Example 59

Preparation of 1,1'-[1,3-cyclohexanebis(methylene)]-[5-(3-trifluoromethylphenyl)biguanide]dihydrochloride

20 The captioned compound was obtained in the same manner as in Example 51 except that p-chloroaniline hydrochloride was replaced by the equimolecular amount of p-trifluoromethylaniline hydrochloride (Yield 2.5 g).

White edged crystals mp. 240~242 °C

Example 60

25

Preparation of 1,1'-[1,3-cyclohexanebis(methylene)]-[5-(2-ethylhexyl)biguanide]dihydrochloride

The captioned compound was obtained in the same manner as in Example 51 except that p-chloroaniline hydrochloride was replaced by the equimolecular amount of p-(2-ethylhexyl)aniline hydrochloride (Yield 4.6 g).

30

White edged crystals mp. 105~107 °C

Example 61

35 Preparation of 1,1'-[1,3-cyclohexanebis(methylene)]-[5-(3,4-dichlorobenzyl)biguanide]dihydrochloride

The captioned compound was obtained in the same manner as in Example 51 except that p-chloroaniline hydrochloride was replaced by the equimolecular amount of 3,4-dichlorobenzylamine hydrochloride (Yield 5.3 g).

40

White edged crystals mp. 231~233 °C

Example 62

Preparation of 1,1'-[1,3-cyclohexanebis(methylene)]-[5-(3,4-dichlorophenyl)biguanide]dihydrochloride

45

The captioned compound was obtained in the same manner as in Example 51 except that p-chloroaniline hydrochloride was replaced by the equimolecular amount of 3,4-dichloroaniline hydrochloride (Yield 3.7 g).

White edged crystals mp. 259~261 °C

50

Example 63

Preparation of 1,1'-[1,3-cyclohexanebis(methylene)]-[5-benzylbiguanide]dihydrochloride

55

The captioned compound was obtained in the same manner as in Example 51 except that p-chloroaniline hydrochloride was replaced by the equimolecular amount of benzylamine hydrochloride (yield 5.9 g).

White edged crystals mp. 208~211 °C

Example 64

Preparation of 1,1'-[1,3-cyclohexanebis(methylene)]-[5-(4-chlorobenzyl)biguanide]dihydrochloride

- 5 The captioned compound was obtained in the same manner as in Example 51 except that p-chloroaniline hydrochloride was replaced by the equimolecular amount of 4-chlorobenzylamine hydrochloride (Yield 4.4 g).

White edged crystals mp. 204~207 ° C

10

Prescription 3 of disinfectant	
Gluconic acid salt of Example 51	5 g
Nonionic surface active agent (Polyoxyethylene phenylether)	3.75 g
Distilled water for injection	proper
Whole volume	100 ml

15

20

Prescription 4 of disinfectant	
Gluconic acid salt of Example 51	0.5 g
Nonionic surface active agent (Polyoxyethylene phenylether)	0.375 g
Ethanol	8.3 ml
Distilled water for injection	proper
Whole volume	100 ml

25

[Antibacterial activity test]

30

To investigate the antibacterial actions of the compounds obtained in Examples 51 to 64, the minimum inhibitory concentration (MIC) was determined by employing the standard method of Japan Society of Chemotherapysame as mentioned above. The results are shown in Table 3.

35

40

45

50

55

Table 3 (1/2)

Strain	MIC $\mu\text{g}/\text{ml}$ (10^4 cells/ml)						
	Examples						
	51	52	53	54	55	56	57
<i>S. aureus</i> FDA 209P	0.39	0.39	0.39	0.78	3.13	3.13	1.56
<i>S. epidermidis</i> ATCC 12228	0.39	0.39	0.39	0.39	3.13	1.56	1.56
<i>E. coli</i> NIHJ JC-2	3.13	6.25	3.13	6.25	25	25	12.5
<i>S. typhi</i> NCTC 8393	1.56	6.25	3.13	6.25	6.25	12.5	6.25
<i>S. marcescens</i> IFO 12648	6.25	12.5	6.25	50	50	50	50
<i>P. aeruginosa</i> ATCC 10145	12.5	100	25	50	100	>100	100
<i>C. albicans</i> IFO 1385	3.13	6.25	6.25	6.25	25	50	100
<i>A. fumigatus</i> IFM 4942	125	250	125	250	400	>400	>400

Table 3 (2/2)


Strain	MIC $\mu\text{g}/\text{ml}$ (10^6 cells/ml)						
	59	Examples					Control agent
		61	62	63	64		
<i>S. aureus</i> FDA 209P	1. 56	3. 13	0. 78	3. 13	1. 56	1. 56	
<i>S. epidermidis</i> ATCC 12228	1. 56	1. 56	0. 39	1. 56	1. 56	0. 78	
<i>E. coli</i> NIHJ JC-2	12. 5	12. 5	6. 25	25	25	1. 56	
<i>S. typhi</i> NCTC 8393	6. 25	12. 5	6. 25	25	6. 25	1. 56	
<i>S. marcescens</i> IFO 12648	25	50	25	50	25	25	
<i>P. aeruginosa</i> ATCC 10145	>100	>100	100	>100	>100	>100	
<i>C. albicans</i> IFO 1385	100	100	6. 25	100	100	5	
<i>A. fumigatus</i> IFM 4942	>200	>200	125	>200	>200	>200	

Claims

1. A biguanide derivative or its salt selected from:

N^1 -(3,4-dichlorobenzyl)- N^5 -octyl-biguanide or its salt,
 N^1 -(4-chlorophenyl)- N^5 -(3,4-dichlorobenzyl)-biguanide or its salt,
 N^1 -(3,4-dichlorophenyl)- N^5 -octyl-biguanide or its salt,
 N^1 -benzyl- N^5 -dodecyl-biguanide or its salt,
 N^1 -benzyl- N^5 -decyl-biguanide or its salt,
 N^1 -(3,4-dichlorobenzyl)- N^5 -(3-trifluoromethylphenyl)-biguanide or its salt,
 N^1 -(3,4-dichlorobenzyl)- N^5 -(4-bromophenyl)-biguanide or its salt,
 N^1 -(3,4-dichlorobenzyl)- N^5 -(4-iodophenyl)-biguanide or its salt,
 N^1 -(3,4-dichlorobenzyl)- N^5 -(2,4-dichlorophenyl)-biguanide or its salt,
 N^1 -(3,4-dichlorobenzyl)- N^5 -(3,4-dichlorophenyl)-biguanide or its salt,


- N¹-(3,4-dichlorobenzyl)-N⁵-(2,3,4-trichlorophenyl)-biguanide or its salt,
 N¹-(3,4-dichlorobenzyl)-N⁵-(3,4-dimethylphenyl)-biguanide or its salt,
 N¹-(3,4-dichlorobenzyl)-N⁵-(3,4-methylenedioxyphenyl)-biguanide or its salt,
 N¹-(3,4-dichlorobenzyl)-N⁵-(4-*t*-butylphenyl)-biguanide or its salt,
 5 N¹-(3,4-dichlorobenzyl)-N⁵-(4-ethylthiophenyl)-biguanide or its salt,
 N¹-(3,4-dichlorobenzyl)-N⁵-(*n*-decyl)-biguanide or its salt,
 N¹-(3,4-dichlorobenzyl)-N⁵-(1,1,3,3-tetramethylbutyl)-biguanide or its salt,
 N¹-(3,4-dichlorobenzyl)-N⁵-hexyl-biguanide or its salt,
 N¹-(3,4-dichlorobenzyl)-N⁵-(2-ethoxyethyl)-biguanide or its salt,
 10 N¹-(3,4-dichlorobenzyl)-N⁵-(2-(2-hydroxyethoxy)ethyl)-biguanide or its salt,
 N¹-(3,4-dichlorobenzyl)-N⁵-(3-diethylaminopropyl)-biguanide or its salt,
 N¹-(3,4-dichlorobenzyl)-N⁵-(3-(2-ethylhexyloxy)-propyl)-biguanide or its salt,
 N¹-(3,4-dichlorobenzyl)-N⁵-[(3-isopropoxy)propyl]-biguanide or its salt,
 N¹-(3,4-dichlorobenzyl)-N⁵-[(2-diethylamino)-ethyl]-biguanide or its salt,
 15 N¹-(3,4-dichlorobenzyl)-N⁵-[(3-butyl)-propyl]-biguanide or its salt,
 N¹-(3,4-dichlorobenzyl)-N⁵-[3(di-*n*-butylamino)propyl]-biguanide or its salt,
 N¹-(4-chlorophenyl)-N⁵-cyclohexylmethyl-biguanide or its salt,
 N¹-(4-chlorophenyl)-N⁵-(3-trifluoromethylphenyl)-biguanide or its salt,
 N¹-(4-chlorophenyl)-N⁵-(4-ethylthiophenyl)-biguanide or its salt,
 20 N¹-(4-chlorophenyl)-N⁵-(4-chlorobenzyl)-biguanide or its salt,
 N¹-(4-chlorophenyl)-N⁵-(2,4-dichlorobenzyl)-biguanide or its salt,
 N¹-(4-chlorophenyl)-N⁵-(4-acetylaminophenyl)-biguanide or its salt,
 N¹-(4-chlorophenyl)-N⁵-(3,4-methylenedioxyphenyl)-biguanide or its salt,
 N¹-(4-chlorophenyl)-N⁵-(3,4-methyleneoxydibenzyl)-biguanide or its salt,
 25 N¹-(4-chlorobenzyl)-N⁵-octyl-biguanide or its salt,
 N¹-(4-chlorobenzyl)-N⁵-(4-chlorobenzyl)-biguanide or its salt,
 N¹-(4-chlorobenzyl)-N⁵-decyl-biguanide or its salt,
 N¹-(4-chlorobenzyl)-N⁵-dodecyl-biguanide or its salt,
 N¹-(4-chlorobenzyl)-N⁵-isobutyl-biguanide or its salt,
 30 N¹-(4-chlorobenzyl)-N⁵-(3,4-dichlorophenyl)-biguanide or its salt,
 N¹-(3,4-dichlorophenyl)-N⁵-(3,4-methylenedioxybenzyl)-biguanide or its salt,
 N¹-(3,4-dichlorophenyl)-N⁵-hexyl-biguanide or its salt,
 N¹-(3,4-dichlorophenyl)-N⁵-decyl-biguanide or its salt,
 35 1,1'-[1,3-cyclohexanebis(methylene)]-bis[5-(4-chlorophenyl)biguanide] or its salt,
 1,1'-[1,3-cyclohexanebis(methylene)]-bis[5-(4-iodophenyl)biguanide] or its salt,
 1,1'-[1,3-cyclohexanebis(methylene)]-bis[5-(4-bromo-phenyl)biguanide] or its salt,
 1,1'-[1,3-cyclohexanebis(methylene)]-bis[5-(2,4-dichlorophenyl)biguanide] or its salt,
 1,1'-[1,3-cyclohexanebis(methylene)]-bis[5-(2,4-difluorophenyl)biguanide] or its salt,
 1,1'-[1,3-cyclohexanebis(methylene)]-bis[5-(4-*t*-butylphenyl)biguanide] or its salt,
 40 1,1'-[1,3-cyclohexanebis(methylene)]-bis[5-(3,4-dimethylphenyl)biguanide] or its salt,
 1,1'-[1,3-cyclohexanebis(methylene)]-bis[5-(4-ethylthiophenyl)biguanide] or its salt,
 1,1'-[1,3-cyclohexanebis(methylene)]-bis[5-(3-trifluoromethylphenyl)biguanide] or its salt,
 1,1'-[1,3-cyclohexanebis(methylene)]-bis[5-(2-ethylhexyl)biguanide] or its salt,
 1,1'-[1,3-cyclohexanebis(methylene)]-bis[5-(3,4-dichlorobenzyl)biguanide] or its salt,
 45 1,1'-[1,3-cyclohexanebis(methylene)]-bis[5-(3,4-dichlorophenyl)biguanide] or its salt,
 1,1'-[1,3-cyclohexanebis(methylene)]-bis[5-benzyl biguanide] or its salt, and
 1,1'-[1,3-cyclohexanebis(methylene)]-bis[5-(4-chlorobenzyl)biguanide] or its salt.
2. A biguanide derivative or its salt selected from:
 50 N¹-(3,4-dichlorobenzyl)-N⁵-octyl-biguanide or its salt,
 N¹-(4-chlorophenyl)-N⁵-(3,4-dichlorobenzyl)-biguanide or its salt,
 N¹-(3,4-dichlorophenyl)-N⁵-octyl-biguanide or its salt, and
 N¹-benzyl-N⁵-dodecyl-biguanide or its salt.
- 55 3. N¹-(3,4-Dichlorobenzyl)-N⁵-octyl-biguanide or its salt.
4. A biguanide derivative or its salt of claim 3, selected from:
 N¹-(3,4-dichlorobenzyl)-N⁵-octyl-biguanide hydrochloride.



- 20 (where R³ represents phenyl group which may possess one to three groups selected from a group consisting of halogen atom, lower alkyl group, halogen substituted lower alkyl group and lower alkylthio group as substituent; phenyl lower alkyl group which may possess halogen atom as substituent on the phenyl ring; or alkyl group, and A denotes a lower alkylene group).
- 25 7. 1,1'-[1,3-Cyclohexanebis(methylene)]-bis[5-(3,4-dichlorobenzyl)biguanide] or its salt.
8. 1,1'-[1,3-Cyclohexanebis(methylene)]-bis[5-(3,4-dichlorophenyl)biguanide] or its salt.
9. A process for preparing a biguanide derivative or its salt, which comprises;
- 30 (i) reacting an amine or its salt expressed by a general formula:




where R¹ represents 3,4-dichlorobenzyl group, 4-chlorophenyl group, 3,4-dichlorophenyl group, benzyl group or 4-chlorobenzyl group with N¹-cyanoguanidine compound expressed by a general formula:



41

group, decyl group, dodecyl group, isobutyl group, 3,4-dichlorophenyl group, or hexyl group to obtain biguanide derivative expressed by a general formula:



(where R¹ and R² are same as defined above) or its salt, selected from N¹-(3,4-dichlorobenzyl)-N⁵-octyl-biguanide or its salt,

N¹-(4-chlorophenyl)-N⁵-(3,4-dichlorobenzyl)-biguanide or its salt,

N¹-(3,4-dichlorophenyl)-N⁵-octyl-biguanide or its salt,

N¹-benzyl-N⁵-dodecyl-biguanide or its salt,

N¹-benzyl-N⁵-decyl-biguanide or its salt,

N¹-(3,4-dichlorobenzyl)-N⁵-(3-trifluoromethylphenyl)-biguanide or its salt,

N¹-(3,4-dichlorobenzyl)-N⁵-(4-bromophenyl)-biguanide or its salt,

N¹-(3,4-dichlorobenzyl)-N⁵-(4-iodophenyl)-biguanide or its salt,

N¹-(3,4-dichlorobenzyl)-N⁵-(2,4-dichlorophenyl)-biguanide or its salt,

N¹-(3,4-dichlorobenzyl)-N⁵-(3,4-dichlorophenyl)-biguanide or its salt,

N¹-(3,4-dichlorobenzyl)-N⁵-(2,3,4-trichlorophenyl)-biguanide or its salt,

N¹-(3,4-dichlorobenzyl)-N⁵-(3,4-dimethylphenyl)-biguanide or its salt,

N¹-(3,4-dichlorobenzyl)-N⁵-(3,4-methylenedioxyphenyl)-biguanide or its salt,

N¹-(3,4-dichlorobenzyl)-N⁵-(4-t-butylphenyl)-biguanide or its salt,

N¹-(3,4-dichlorobenzyl)-N⁵-(4-ethylthiophenyl)-biguanide or its salt,

N¹-(3,4-dichlorobenzyl)-N⁵-(n-decyl)-biguanide or its salt,

N¹-(3,4-dichlorobenzyl)-N⁵-(1,1,3,3-tetramethylbutyl)-biguanide or its salt,

N¹-(3,4-dichlorobenzyl)-N⁵-hexyl-biguanide or its salt,

N¹-(3,4-dichlorobenzyl)-N⁵-(2-ethoxyethyl)-biguanide or its salt,

N¹-(3,4-dichlorobenzyl)-N⁵-(2-(2-hydroxyethoxy) ethyl)-biguanide or its salt,

N¹-(3,4-dichlorobenzyl)-N⁵-(3-diethylaminopropyl)-biguanide or its salt,

N¹-(3,4-dichlorobenzyl)-N⁵-(3-(2-ethylhexyloxy)-propyl)-biguanide or its salt,

N¹-(3,4-dichlorobenzyl)-N⁵-[(3-isopropoxy)propyl]-biguanide or its salt,

N¹-(3,4-dichlorobenzyl)-N⁵-[(2-diethylamino)-ethyl]-biguanide or its salt,

N¹-(3,4-dichlorobenzyl)-N⁵-[(3-butyl)-propyl]-biguanide or its salt,

N¹-(3,4-dichlorobenzyl)-N⁵-[3(di-n-butylamino)propyl]-biguanide or its salt,

N¹-(4-chlorophenyl)-N⁵-cyclohexylmethyl-biguanide or its salt,

N¹-(4-chlorophenyl)-N⁵-(3-trifluoromethylphenyl)-biguanide or its salt,

N¹-(4-chlorophenyl)-N⁵-(4-ethylthiophenyl)-biguanide or its salt,

N¹-(4-chlorophenyl)-N⁵-(4-chlorobenzyl)-biguanide or its salt,

N¹-(4-chlorophenyl)-N⁵-(2,4-dichlorobenzyl)-biguanide or its salt,

N¹-(4-chlorophenyl)-N⁵-(4-acetylaminophenyl)-biguanide or its salt,

N¹-(4-chlorophenyl)-N⁵-(3,4-methylenedioxyphenyl)-biguanide or its salt,

N¹-(4-chlorophenyl)-N⁵-(3,4-methyleneoxydibenzyl)-biguanide or its salt,

N¹-(4-chlorobenzyl)-N⁵-octyl-biguanide or its salt,

N¹-(4-chlorobenzyl)-N⁵-(4-chlorobenzyl)-biguanide or its salt,

N¹-(4-chlorobenzyl)-N⁵-decyl-biguanide or its salt,

N¹-(4-chlorobenzyl)-N⁵-dodecyl-biguanide or its salt,

N¹-(4-chlorobenzyl)-N⁵-isobutyl-biguanide or its salt,


N¹-(4-chlorobenzyl)-N⁵-(3,4-dichlorophenyl)-biguanide or its salt,

N¹-(3,4-dichlorophenyl)-N⁵-(3,4-methylenedioxybenzyl)-biguanide or its salt,


N¹-(3,4-dichlorophenyl)-N⁵-hexyl-biguanide or its salt,

N¹-(3,4-dichlorophenyl)-N⁵-decyl-biguanide or its salt;

(II) N⁵-cyanoguanidine compound expressed by a general formula:



(wherein R¹ is same as defined above), is reacted with and an amine or its salt expressed by general a formula:




(wherein R² is same as defined above) to obtain biguanide derivative expressed by the above formula (1) or its salt;


(III) reacting N¹-cyano-N³-octyl-guanidine with 3,4-dichlorobenzylamine or its acid-addition salt, or reacting N¹-(3,4-dichlorobenzyl)-N³-cyano-guanidine with octylamine or its acid-addition salt to obtain N¹-(3,4-dichlorobenzyl)-N⁵-octyl biguanide or its salt;

(IV) reacting N¹-cyano-N³-(3,4-dichlorobenzyl)-guanidine with 4-chloroaniline or its acid-addition salt, or reacting N¹-(4-chlorophenyl)-N³-cyano-guanidine with 3,4-dichlorobenzylamine or its acid-addition salt, to obtain N¹-(4-chlorophenyl)-N⁵-(3,4-dichlorobenzyl)-biguanide or its salt;


(V) reacting 1,1'-[1,3-cyanohexanebis(alkylene)]-bis(3-cyanoguanidine) expressed by a formula:




(where A denotes a lower alkylene group), with an amine or its salt expressed by a formula:




(where R³ represents phenyl group which may possess one to three groups selected from a group consisting of halogen atom, lower alkyl group, halogen substituted lower alkyl group and lower alkylthio group as substituent, phenyl lower alkyl group which may possess halogen atom as substituent on phenyl ring, or alkyl group) to obtain biguanide derivative or its salt of claim 6 expressed by a formula (2):



(where A and R^3 are same as defined in claim 6); and
(VI) reacting a diamine or its salt expressed by a formula:



(where A is same as defined above) with a cyanoguanidine compound expressed by a formula:



(where R^3 is same as defined above) to obtain biguanide derivative or its salt of claim 6.

10. A disinfectant composition comprising an effective amount of a biguanide derivative or its salt of claim 1, and its carrier.
11. A disinfectant composition comprising an effective amount of biguanide derivative or its salt of claim 2, and its carrier.
12. A disinfectant composition comprising an effective amount of a biguanide derivative or its salt of claim 3, and its carrier.
13. A disinfectant composition comprising an effective amount of a biguanide derivative or its salt of claim 5, and its carrier.
14. A disinfectant composition comprising an effective amount of a biguanide derivative or its salt of claim 6, and its carrier.
15. A disinfectant composition comprising an effective amount of a biguanide derivative or its salt of claim 7 or 8, and its carrier.
16. Use of biguanide derivative or salt of claim 1, for manufacture of disinfectant for humans, animals, or medical appliances.
17. Use of biguanide derivative or salt of claim 2, for manufacture of disinfectant for humans, animals, or medical appliances.
18. Use of biguanide derivative or salt of claim 3 or 5, for manufacture of disinfectant for humans, animals, or medical appliances.
19. Use of biguanide derivative or salt of claim 6, for manufacture of disinfectant for humans, animals, or

medical appliances.

5

10

15

20

25

30

35

40

45

50

55

(19)



Europäisches Patentamt

Europ an Patent Office

Office européen des brevets



(11) Publication number:

0 507 317 A3

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 92105776.6

(51) Int. Cl.⁵: **C07C 279/26, A61K 31/155**

(22) Date of filing: 03.04.92

(30) Priority: 05.04.91 JP 73202/91
 04.09.91 JP 224306/91
 19.06.91 JP 147644/91

(43) Date of publication of application:
 07.10.92 Bulletin 92/41

(84) Designated Contracting States:
**AT BE CH DE DK ES FR GB GR IT LI LU MC
 NL PT SE**

(88) Date of deferred publication of the search report:
 24.02.93 Bulletin 93/08

(71) Applicant: OTSUKA PHARMACEUTICAL CO.,
LTD.
 9, Kandatsukasacho 2-chome
 Chiyoda-ku Tokyo 101(JP)

(72) Inventor: Ishikawa, Hiroshi
 18-14, Seifu-cho

Otsu-shi, Shiga 520-02(JP)

Inventor: Yasumura, Koichi

8-11-401, Uchidehama**Otsu-shi, Shiga 520(JP)**

Inventor: Tsubouchi, Hidetsugu

2-6-608, Nionohama 2-chome**Otsu-shi, Shiga 520(JP)**

Inventor: Higuchi, Yukio

8-17, Chodo 2-chome**Higashiosaka-shi, Osaka 577(JP)**


Inventor: Tamaoka, Hisashi


5-32, Aza-ekogawa, Shinkirai, Kitajima-cho**Itano-gun, Tokushima 771-02(JP)**

(74) Representative: Hansen, Bernd, Dr.

Dipl.-Chem. et al**Hoffmann, Eitle & Partner Patent- und****Rechtsanwälte Arabellastrasse 4 Postfach****81 04 20****W-8000 München 81 (DE)**(54) **Biguanide derivatives, manufacturing method thereof, and disinfectants containing the derivatives.**

(57) The invention presents a biguanide derivative or its salt expressed by a formula:

(where R¹ and R² are as defined in Specification), or formula:**EP 0 507 317 A3**



(2)

(where A and R³ is as defined in specification). This biguanide derivative or its salt is preferably used as the effective amount of a disinfectant for humans, animals, medical appliances, etc.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 92 10 5776

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
X	US-A-3 183 230 (S.L. SHAPIRO ET AL.) * column 1; claims * ---	1,9,10	C07C279/26 A61K31/155
X	US-A-4 183 958 (M.J. BROWN ET AL.) * claims; examples * ---	1,9,10	
X	FR-M-2 100 (U.S. VITAMIN & PHARMACEUTICAL CORPORATION) * column 1; claims * ---	1,9,10	
X	FR-A-1 015 134 (IMPERIAL CHEMICAL INDUSTRIES PLC) * claims * ---	1,9	
X	FR-A-2 085 665 (FARBENFABRIKEN BAYER AKTIEN GESELLSCHAFT) * claims * ---	1,9	
X	US-B-403 243 (J. DIAMOND ET AL.) * examples * ---	1,9	
A,D	GB-A-1 095 902 (STERLING DRUG INC.) * claims * ---	1,6-9, 10, 14-16,19	TECHNICAL FIELDS SEARCHED (Int. Cl.5) C07C
A	US-A-4 567 174 (P.N. EDWARDS ET AL.) * claims * -----	1,6-9, 10, 14-16,19	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 10 DECEMBER 1992	Examiner SANCHEZ GARCIA J.M.
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- @ : member of the same patent family, corresponding document			

EPO FORM 1503 01.92 (P0401)

